



Bio-accumulation or bio-concentration of radionucleotides through food chain

Concentration factor	Concentration in fish body Concentration in sea water
Materials	Concentration Factor of marine fish
Cs	5 ~ 100
I	10
U	10
Pt	3.5
Hg	360 ~ 600
DDT	12000
PCB	1200 ~ 1000000



Very low Concentration Factors

Reference:
Fujio Kasamatsu
bio-concentration Edit. N. Yamagata ,
Radioisotopes 48, 1999.



- Bio-accumulation or bio-concentration of radionucleotides through food chain is not increasing.

Why are not accumulated ?



Iodine and Cesium

- Iodinesolid/gaseous (sublimation nucleotide)

I-131 (Half life time: 8.04 days)

- Cs.....solid , behaves like potassium :

does not accumulate to specific organs


Cs-137 (Half life time :30.1years),

Cs-134 (Half life time: 2.07years)

Periodic table

	1A	2A	3A	4A	5A	6A	7A	8	1B	2B	3B	4B	5B	6B	7B	0					
1	H																				He
2	Li	Be																			
3	Na	Mg																			
4	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr			
5	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe			
6	Cs	Ba	*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn			
7	Fr	Ra	**																		
*	La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu						
**	Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr						


The flow of salts in marine fish body



- Radionucleotides excrete, not accumulate.
- The concentration in fish is depend on the concentration of environmental water .


(Ref: Fundamental physiology of fish
Edit. K. Aida)

Comparison of Cs-137 concentration between sea water and fish body




- Fish body concentration depends on sea water concentration

Excretion of radio nucleotides



Biological half time of

$\text{Cs-137}=50 \text{ days}$



The half of Cs-137 is excreted in 50days. (Laboratory work result)

- In natural condition Cs-137 excretes quickly.

Ref:

K. Yoshida , JCAC 34, 1999.
F. Kasamatsu, Radioisotopes 48, 1999.